

May 1996

SUPERLUMINAL MATTER AND HIGH-ENERGY COSMIC RAYS

L. GONZALEZ-MESTRES

Laboratoire de Physique Corpusculaire, Collège de France,
11 pl. Marcellin-Berthelot, 75231 Paris Cedex 05 , France

High-energy cosmic ray events present important challenges to particle astrophysics. Their nature and origin are often not well understood and, as they occur in an energy domain not accessible to particle accelerators, there is no clear guide to their interpretation. We recently proposed [1] a new, non-tachyonic, possible class of superluminal particles which may play an important cosmological role [2]. In this note, we would like to point out that such particles, if they exist, are able to produce very high energy cosmic rays compatible with unexplained discoveries reported in the literature.

1. THE NEW SUPERLUMINAL PARTICLES

Since we know from particle physics that vacuum is not an empty entity but a material medium with many internal degrees of freedom, there is no fundamental reason to interpret Lorentz invariance as an absolute property of space and time previous to the existence of matter. Instead, we may consider it as an approximate dynamical property of the equations describing an important sector of matter (the "ordinary" sector, i.e. the particles whose critical speed in vacuum is the speed of light) above a certain distance scale (k_0^{-1} , where k_0 is a critical wave vector scale) related to a fundamental phase transition (at $T \approx k^{-1} \hbar c k_0$, k = Boltzmann constant, $\hbar$ = Plack constant [2]). Then, ordinary Lorentz invariance would not necessarily be universal and other degrees of freedom can exist leading to particles with different critical speeds in vacuum (the "superluminal" sectors, with critical speeds $c_i \gg c$ where the subscript i stands for the i -th superluminal sector). If m is the inertial mass, a particle belonging to the i -th superluminal sector would have [1] rest energy:

$$E_{rest} = m c_i^2 \quad (1)$$

leading potentially to very large rest energies. Annihilation of pairs of superluminal particles into ordinary ones can therefore release very large kinetic energies and provide a new source of high-energy cosmic rays. If superluminal matter exists, it is likely to have dominated the evolution of the very early Universe [2] and be still present playing a crucial role at large scales. The study of fundamental cosmological problems (e.g. nature of the big bang, horizon problem, cosmological constant...) would be considerably modified.

Gravitational coupling of superluminal particles to ordinary ones is expected to be weak. Assuming that each superluminal sector has its own Lorentz metric $g_{[i]\mu\nu}$ ($[i]$ for the i -th sector), with c_i setting the speed scale, we may expect each sector to generate its own gravity with a coupling constant κ_i and a new sectorial graviton. Gravitation would be a single and universal interaction in the limit where all c_i tend to c and where a single metric can be used, but not in other situations. As an ansatz, we can assume that the static gravitational coupling between two different sectors is lowered by a factor proportional to a positive power of the ratio between the two critical speeds (the smallest speed divided by the largest one). Static gravitational forces between ordinary matter and matter of the i -th superluminal sector would then be proportional to a positive power of c/c_i which can be a very small number. Experiment seems to suggest that other known gauge interactions (e.g. electroweak, strong) also couple weakly to superluminal matter and that, if such matter exists, it can be produced only at very high energies (future generations of accelerators?). Interactions between ordinary and superluminal matter will break ordinary Lorentz invariance. This will happen basically at very high energy and very short distance, and is not incompatible with the success of conventional tests of relativity.

2. THE RELEVANCE OF COSMIC RAY STUDIES

Because of the weak coupling to ordinary particles, concentrations of superluminal matter would not necessarily follow the same pattern as "ordinary" galaxies and clusters of galaxies (made of ordinary matter), nor would they need to occur at the same places. Although we expect correlations between the distributions of ordinary and superluminal matter at large scales, it may be difficult to reasonably assume that the gravitational role of galactic halos be due to superluminal particles. Superluminal particles can be very abundant and even provide most of the dark matter at cosmic scale, but they will be extremely difficult to detect if they interact very weakly with ordinary matter. Similarly, astrophysical objects made of superluminal matter may elude all conventional observational techniques and be extremely difficult to find. Cosmic rays can then play a crucial role.

a) Ordinary primaries. *Annihilation* of pairs of slow superluminal particles *into ordinary particles*, releasing very high kinetic energies from the superluminal rest masses, would yield a unique *cosmic signature* allowing cosmic ray detectors to search for this new kind of matter in the present Universe. *Collisions* of high-energy superluminal particles *with extra-terrestrial ordinary matter* may also yield high-energy ordinary cosmic rays. High-energy superluminal particles can be produced from acceleration, decays, explosions... in astrophysical objects made of superluminal matter. Pairs of slow superluminal particles can also annihilate into particles of another superluminal sector with lower c_i , converting most of the rest energies into a large amount of kinetic energy. Superluminal particles moving at $v > c$ will release "*Cherenkov*" *radiation in vacuum*, i.e. spontaneous emission of particles of a lower critical speed c_i (for $v > c_i$) including ordinary ones, providing a new source of (superluminal or ordinary) high-energy cosmic rays.

b) Superluminal primaries. High-energy superluminal particles can directly *reach the earth* and undergo collisions inside the atmosphere, producing many secondaries like

ordinary cosmic ray events. They can also interact with the rock or with water near some underground or underwater detector, coming from the atmosphere or after having crossed the earth, and producing clear signatures. Contrary to neutrinos, whose flux is strongly attenuated by the earth at energies larger than 10^6 GeV, superluminal particles will in principle not be stopped by earth at these energies. Even if the interaction probability is expected to be very weak, the rate for superluminal cosmic ray events can be observable if we are surrounded by important concentrations of superluminal matter.

The possibility that superluminal matter exists, and that it plays nowadays an important role in our Universe, should be kept in mind when addressing the two basic questions raised by the analysis of cosmic ray events: a) the nature of the cosmic ray primary; b) the identification of the source of the cosmic ray. If the primary is a superluminal particle, it will escape conventional criteria for particle identification and most likely produce a specific signature. If the source is superluminal, it can be located anywhere (and even be a free particle) and will not be at the same place as conventional sources of ordinary cosmic rays. High-energy cosmic ray events originating from superluminal sources will for the first time provide hints on the location of such sources. The energy dependence of the events should also be taken into account. At very high energies, the Greisen-Zatsepin-Kuzmin cut-off [3] does not in principle hold for cosmic ray events originating from superluminal matter: this is obvious if the primaries are superluminal particles that we expect to interact very weakly with the cosmic microwave background, but is also true for ordinary primaries as we do not expect them to be produced at the locations of ordinary sources and there is no upper bound to their energy around 100 EeV. To date, there is no well established interpretation [3, 4] of the highest-energy cosmic ray events [5]. Primaries (ordinary or superluminal) originating from superluminal particles are acceptable candidates and naturally escape several problems faced by cosmic rays produced at ordinary sources.

References

- [1] L. Gonzalez-Mestres, "Properties of a possible class of particles able to travel faster than light", Proceedings of the Moriond Workshop on "Dark Matter in Cosmology, Clocks and Tests of Fundamental Laws", Villars (Switzerland), January 21 - 28 1995, Ed. Frontières. Paper astro-ph/9505117 of electronic library.
- [2] L. Gonzalez-Mestres, "Cosmological implications of a possible class of particles able to travel faster than light", Proceedings of the IV International Conference on Theoretical and Phenomenological Aspects of Underground Physics, TAUP95, Toledo September 1995, Ed. Nuclear Physics Proceedings. Paper astro-ph/9601090 of electronic library.
- [3] K. Greisen, Phys. Rev. Lett. 16, 748 (1966); G.T. Zatsepin and V. A. Kuzmin, Pisma Zh. Eksp. Teor. Fiz. 4, 114 (1966).
- [4] F. W. Stecker, Phys. Rev. 180, 1264 (1969); F. Halzen, R.A. Vazquez, T. Stanev and V.P. Vankov, Astropart. Phys. 3, 151 (1995); T.J. Weiler and T.W. Kephart, paper presented at the Dark Matter'96 Conference, UCLA February 14-16 1996, astro-ph/9605156.
- [5] S. Yoshida et al., Astropart. Phys. 3, 105 (1995); D.J. Bird et al. Astrophys. J. 424, 491 (1994) and Astrophys. J. 441, 144 (1995); M.A. Lawrence, R.J.O. Reid and A.A. Watson, J. Phys. G17, 733 (1991); B.N. Afanasiev et al. in Proceedings of the XXIV International Cosmic Ray Conference, Rome 1995, Vol. 2, page 756.